

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis	
1. FRUITS FRESH OR FROZEN; NUTS																				
i) Citrus fruit																				
Grapefruit (Shaddocks, pomelos, sweeties, tangelo, ugli and other hybrids)	3	875	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Grapefruit (Shaddocks, pomelos, sweeties, tangelo, ugli and other hybrids) (China, Turkey, South Africa)	5	480	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Oranges	1	2500	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Lemons	1	2500	x					O*	O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample. *Recommendation: Overseas goods: ethephon at the beginning of the season Obligation: Sampling of each 4th sample after all process steps ²
Lemons (Turkey)	6	340	x					O*	O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Limes	5	480	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Mandarins (clementine, tangerine and other hybrids)	1	2500	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Mandarins (clementine, tangerine and other hybrids) (Third countries)	3	875	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Other citrus fruits	6	340	x						O(2,4-D)	x	O	O	O							Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
ii) Tree nuts and seeds (shelled or unshelled)																				
Almonds	1	2500	x							x										
Brazil nut	1	2500	x							x										
Cashew nut	1	2500	x							x										
Chestnut	1	2500	x							x										
Chestnut (China, Turkey)	6	340	x							x										Obligation: Sampling after all process steps
Coconut	1	2500	x							x										
Hazelnut	2	1600	x							x										
Macadamia nut	1	2500	x							x										
Pecans	1	2500	x							x										
Pine nuts	1	2500	x							x										
Pistachio	1	2500	x							x										
Walnut	2	1600	x							x										
Peanut	2	1600	x							x				x*						*Obligation: for each 3rd sample
Other nuts and seeds (shelled or unshelled)	6	340	x							x										
iii) Pome fruit																				
Apple	1	2500	x				x			x										
Apple (Third countries except New Zealand)	3	875	x							x										
Pear	3	875	x			x*	x			x										
Pear (Belgium)	2	1600	x			x*	x			x										*Obligation: for each 10th sample
Pear (Germany, Netherlands, Austria)	1	2500	x			x*	x			x										
Quince	6	340	x							x										
Loquat	1	2500	x							x										
Medlar	1	2500	x							x										
Nashi pear	6	340	x			x*	x			x										*Obligation: for each 10th sample
Other pome fruits	6	340	x							x										
iv) Stone fruits																				
Apricot	2	1600	x	O						x										
Apricot (Turkey)	6	340	x	O						x										
Sweet cherry	4	640	x	O						x										

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloroquat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis	
Sweet cherry (Turkey)	7	170	x	O			x			x				x						
Sour cherry	4	640	x	O			x	x*												*Obligation: for industrial cultivation
Nectarine	1	2500	x	O						x										
Plum (Damson, greengage, mirabelle)	5	480	x	O			x	x**						x						**Obligation: for industrial cultivation
Peach	1	2500	x	O						x										
Other stone fruits	7	170	x	O			x			x				x						
v) Berries and small fruit																				
a) Grapes																				
Table grapes green	1	2500	x							x										
Table grapes green (Egypt, Brazil, India, Namibia, Peru)	2	1600	x			x*				x										*Obligation: for origin India
Table grapes green (Turkey)	7	170	x							x										
Table grapes blue	2	1600	x			x*		x		x										*Obligation: for origin India
Kiwiberry (mini kiwi)	2	1600	x							x										
b) Strawberries																				
Strawberry (Outdoor)	1	2500	x							x										
Strawberry (Outdoor) (North Africa)	2	1600	x							x										
Strawberry (Greenhouse)	2	1600	x							x										
Strawberry (Greenhouse) (Netherlands)	1	2500	x							x										
c) Cane fruit																				
Blackberry	6	340	x							x										
Blackberry (Mexico)	7	170	x							x										
Raspberry	3	875	x							x										
Dewberry (Loganberry)	6	340	x							x				x*						*Obligation: for each 3rd sample
Other cane fruits	7	170	x							x										
d) Other small fruit and berries																				
Cultivated Blueberry	3	875	x							x										
Cowberry	5	480	x							x										
Cranberry	2	1600	x							x				x						
Currant (red, black and white)	6	340	x							x										
Gooseberry	4	640	x							x										
Jostaberry	3	875	x							x										
Rose hip	3	875	x							x										
Mulberry	3	875	x							x				x						
Elderberries (wild rowan berry)	3	875	x							x										
Cape gooseberry; Physalis	2	1600	x							x										
Other small fruits and berries	6	340	x							x				x						
vi) Miscellaneous fruit																				
a) Edible peel																				
Date	1	2500	x							x				x*						*Obligation: for each 3rd sample
Fig	1	2500	x					x**		x				x*						*Obligation: for each 3rd sample **Obligation: for origin Brazil, for origin Turkey at the beginning of the season until end of August
Table olive	5	480	x							x										
Kumquat	4	640	x					x**		x				x*						*Obligation: for each 3rd sample **Obligation: at the beginning of the season
Carambola	6	340	x							x				x*						*Obligation: for each 3rd sample
Kaki; Japanese persimmons	4	640	x					x*		x										*Obligation: for origin Spain at the beginning of the season
Other miscellaneous fruits with edible peel	6	340	x							x										
b) Inedible peel, small																				
Kiwi	1	2500	x							x										
Lychee	4	640	x							x				x	x					
Maracuja; Passionfruit (Granadilla)	7	170	x							x				x						
Maracuja; Passionfruit (Granadilla) (Columbia)	9	45	x							x				x						

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis	
Prickly pear; cactus fruit; pitaya	9	45	x							x										
Other small miscellaneous fruits with inedible peel	9	45	x							x										
c) Inedible peel, large																				
Avocado	5	480	x							x										
Avocado (Chile, Peru, Columbia)	7	170	x							x						x*				*Obligation: Cadmium
Banana	1	2500	x							x										
Mango	4	640	x							x										
Papaya; Tamarillo	8	90	x							x										
Pomegranate	5	480	x							x										
Pomegranate (Turkey)	7	170	x							x										
Pineapple	1	2500	x					x		x										
Bread fruit; Jackfruit	6	340	x							x										
Cherimoya; Rambutan	6	340	x							x										
Durian	6	340	x							x										
Guava	6	340	x							x										
Other large miscellaneous fruits with inedible peel	8	90	x							x										

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis
2. VEGETABLES FRESH OR FROZEN																			
i) Root and tuber vegetables																			
a) Potatoes																			
Potato	1	2500	x							x									
b) Tropical root and tuber vegetables																			
Ginger	7	170	x							x									
Cassava (Dasheen, eddoe (Japanese taro), tannia, manioc)	4	640	x							x									
Sweet potato	1	2500	x							x				x*					*Obligation: for each 3rd sample
Sweet potato (Egypt)	4	640	x		x					x				x*					*Obligation: for each 3rd sample
Yams (Potato bean (yam bean), Mexican yam bean)	4	640	x							x				x*					*Obligation: for each 3rd sample
Other tropical root and tuber vegetables	7	170	x							x									
(c) Other root and tuber vegetables except sugar beet																			
Beetroot	1	2500	x							x				x*					*Obligation: for each 3rd sample
Carrot (bunch)	1	2500	x							x				x*					*Obligation: for each 3rd sample
Carrot (bulk goods)	1	2500	x							x				x*					*Obligation: for each 3rd sample
Celeriac	4	640	x							x						x			
Horseradish	1	2500	x							x						O			
Jerusalem artichoke	1	2500	x							x				x*					*Obligation: for each 3rd sample
Parsnip	2	1600	x							x				x*		O			*Obligation: for each 3rd sample
Parsley root	2	1600	x							x				x*					*Obligation: for each 3rd sample
Radish (Outdoor)	1	2500	x							x									
Radish (Greenhouse)	1	2500	x							x									
Small radish (Outdoor)	2	1600	x							x									
Small Radish (Greenhouse)	1	2500	x							x									
Scorzonera	1	2500	x							x						x			
White Turnip; Turnip; Swedes	4	640	x							x				x*					*Obligation: for each 3rd sample
Other root and tuber vegetables except sugar beet	4	640	x							x									
ii) Bulb vegetables																			
Garlic	1	2500	x							x									
Onions (Silver onions)	1	2500	x							x									
Onions	1	2500	x							x									
Shallots	1	2500	x							x									
Spring onion	1	2500	x							x									
Other bulb vegetables	1	2500	x							x									

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis
iii) Fruiting vegetables																			
a) Solanaceae																			
Tomato	1	2500	x							x									
Tomato (Turkey)	2	1600	x							x									
Peppers	2	1600	x							x									
Peppers (Germany, Spain)	1	2500	x					x*		x									*Obligation: for origin Greece, Spain, Third countries for the winter season (not valid for green peppers)
Peppers (Turkey)	6	340	x					x*		x									
Chilli peppers	1	2500	x							x									
Chilli peppers (Third Countries)	7	170	x							x									
Aubergines	1	2500	x							x									
Aubergines (Third countries except Morocco)	4	640	x							x									
Okra; Lady's fingers	9	45	x							x				x*					*Obligation: for each 3rd sample
Other Solanaceae	9	45	x							x									
b) Cucurbits- edible peel																			
Cucumber	1	2500	x							x									
Gherkin	1	2500	x							x									
Courgette	3	875	x							x									
Courgette (Netherlands)	1	2500	x							x									
Courgette (Germany, Spain)	2	1600	x							x									
Other cucurbits with edible peel	3	875	x							x									
c) Cucurbits - inedible peel																			
Melon (Muskmelon, Kiwano)	2	1600	x							x									
Squashes (e.g. Spaghetti squash, Halloween squash, Pattypan squash)	2	1600	x							x									
Musky gourd (e.g. Butternut)	2	1600	x							x									
Pumpkin (giant) (e.g. Hokkaido)	2	1600	x							x									
Calabash (e.g. Calabash)	2	1600	x							x									
Pumpkin (others)	2	1600	x							x									
Water melon	1	2500	x							x									
Other cucurbits with inedible peel	2	1600	x							x									
iv) Brassica vegetables																			
a) Flowering brassica																			
Broccoli	4	640	x							x									
Cauliflower (romanesco)	2	1600	x							x									
Cauliflower (romanesco) (Germany)	3	875	x							x									
Other flowering brassica	4	640	x							x									
b) Head brassica																			
Brussels sprout; cabbage sprouts	2	1600	x							x									
Red cabbage	1	2500	x							x									
White cabbage	1	2500	x							x									
Pointed cabbage	1	2500	x							x									
Savoy cabbage	4	640	x							x									
Other head brassica	4	640	x							x									
c) Leafy brassica																			
Chinese cabbage (Indian (Chinese) mustard, pak choy)	2	1600	x							x									
Kale	7	170	x							x									
Other leafy brassica	7	170	x							x									
d) Kohlrabi																			
Kohlrabi (Outdoor) without leaves	2	1600	x							x									
Kohlrabi (Outdoor) with leaves	4	640	x							x									Obligation: Analysis of tuber and leaf
Kohlrabi (Greenhouse) without leaves	2	1600	x							x									
Kohlrabi (Greenhouse) with leaves	4	640	x							x									Obligation: Analysis of tuber and leaf
v) Leaf vegetables and fresh herbs																			
a) Lettuce and other salad plants including Brassicaceae																			
Lamb's lettuce (Outdoor)	3	875	x	O						x									
Lamb's lettuce (Greenhouse)	5	480	x	O						x									
Head lettuce (Outdoor)	2	1600	x	x						x									
Head lettuce (Greenhouse)	2	1600	x	x						x									

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis	
Iceberg lettuce (Outdoor)	1	2500	x	O	x					x										
Coloured lettuce (Lollo, Leaf-oak, Batavia) (Outdoor)	4	640	x	x	x					x										
Coloured lettuce (Lollo, Leaf-oak, Batavia) (Outdoor) (Spain, Netherlands)	3	875	x	x	x					x										
Coloured lettuce (Lollo, Leaf-oak, Batavia) (Greenhouse)	3	875	x	x	x					x										
Romaine lettuce (Outdoor)	1	2500	x	x	x					x										
Escarole/broad-leaf endive (wild chicory, red-leaved chicory, radicchio, curly leaf endive, sugar loaf)	3	875	x	x						x										
Land cress	1	2500	x							x										
Rocket, Rucola	6	340	x		x					x										
Mizuna (Leaves and sprouts of Brassica spp)	6	340	x							x										
Other lettuce and other salad plants including Brassicaceae	6	340	x		x					x										
b) Spinach and similar leaves																				
Spinach	5	480	x	x	x					x							O			
Spinach (Industrial production)	1	2500	x	x	x					x							O			
Purslane (Winter purslane (miner's lettuce))	7	170	x							x				x*						*Obligation: for each 3rd sample
Chard	2	1600	x							x										
Turnip greens	2	1600	x							x										
Other spinach and similar leaves	7	170	x							x										
c) Baby leaf																				
Baby Leaf	2	1600	x	x*						x							O			*Obligation: if not contain brassicaceae
d) Vine leaves (grape leaves)																				
Vine leaves	9	45	x	x						x				x*						*Obligation: for each 3rd sample
e) Water cress																				
Water cress (Water convolvulus, Water clovers, Water mimosas)	4	640	x							x				x*						*Obligation: for each 3rd sample
Water cress (Water convolvulus, Water clovers, Water mimosas) (Third Countries)	7	170	x							x				x*						
f) Witloof																				
Witloof	2	1600	x							x										
g) Fresh herbs and edible flowers⁽⁴⁾																				
Pot herbs																				
⁽⁴⁾ for entire g) fresh herbs (Italy, Third Countries)	9	45	x							x										
Chervil	4	640	x							x										
Chives	4	640	x							x										
Dill leaves	4	640	x							x										
Celery leaves; Sorrel	4	640	x							x										
Coriander leaves	4	640	x							x										
Lovage	4	640	x							x										
Parsley	4	640	x							x										
Sage	6	340	x							x										
Rosemary	6	340	x							x										
Thyme	6	340	x							x										
Basil	4	640	x							x										
Mint	4	640	x							x										
Tarragon (Hyssop)	6	340	x							x										
Wild garlic	4	640	x							x										
Caraway	4	640	x							x										
Bay leaves	6	340	x							x										
Marjoram	4	640	x							x										
Oregano	6	340	x							x										
Savory	4	640	x							x										
Common balm; Lemon balm	4	640	x							x										
Edible flowers	4	640	x							x										
Other fresh herbs and edible flowers	6	340	x							x										
Cut herbs																				

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis	
⁽⁴⁾ for entire q) fresh herbs	9	45	x	x*						x										*Obligation: from November to March for each 4th sample if origin Italy/ Third Countries
Chervil	7	170	x	x*						x										*Obligation: from November to March for each 4th sample
Chives	7	170	x	x*						x										*Obligation: from November to March for each 4th sample
Dill leaves	7	170	x	x*						x										*Obligation: from November to March for each 4th sample
Celery leaves; Sorrel	7	170	x	x*						x										*Obligation: from November to March for each 4th sample
Coriander leaves	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Lovage	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Parsley	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Sage	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Rosemary	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Thyme	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Basil	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Mint	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Tarragon (Hyssop)	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Wild garlic	7	170	x	x*						x										
Caraway	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Bay leaves	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Marjoram	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Oregano	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Savory	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Common balm; Lemon balm	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Edible flowers	7	170	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Other fresh herbs and edible flowers	8	90	x	x*						x										*Obligation: for origin Southern Europe from November to March for each 4th sample
vi) Legume vegetables (fresh)																				
Beans (with pods)	3	875	x							x										
Beans (with pods) (Third Countries)	7	170	x							x										
Beans (without pods)	2	1600	x							x										
Beans (without pods) (Third Countries)	4	640	x							x										*Obligation: for each 3rd sample
Peas (with pods) (Mangetout (sugar peas))	3	875	x							x										*Obligation: for each 3rd sample
Peas (with pods) (Mangetout (sugar peas)) (Third Countries)	9	45	x							x										*Obligation: for each 3rd sample
Peas (without pods)	2	1600	x							x										*Obligation: for each 3rd sample
Peas (without pods) (Third Countries)	4	640	x							x										*Obligation: for each 3rd sample
Other legume vegetables (fresh)	9	45	x							x										
vii) Stem vegetables (fresh)																				
Asparagus white	1	2500	x							x										*Obligation: for each 3rd sample
Asparagus green	1	2500	x							x										
Celery	5	480	x							x										*Obligation: for each 3rd sample
Fennel	1	2500	x							x										*Obligation: for each 3rd sample
Globe artichoke	1	2500	x							x										
Leek	3	875	x							x										
Rhubarb	1	2500	x							x										*Obligation: for each 3rd sample

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce ⁴	Multimethods	Dithiocarbamates	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead, Copper)	Glyphosate	Diquat / Paraquat	Additional analysis
Bamboo shoots	1	2500	x	O						x				x*					*Obligation: for each 3rd sample
Glassworts	7	170	x											x*					*Obligation: for each 3rd sample
Other stem vegetables (fresh)	7	170	x							x									
viii) Fungi																			
Cultivated fungi (common mushroom, oyster mushroom, shi-take)	2	1600	x			x				x				x*				x	*Obligation: for each 3rd sample
Wild fungi (morels, chanterelle)	6	340	x							x				x*		O**			*Obligation: Analysis on radiation exposure (Cäsium-137) *Obligation: for each 3rd sample **Recommendation: Mercury, Cadmium
Other cultivated fungi	6	340	x							x									
3. PULSES, DRY																			
Beans	2	1600	x							x				x*				x	*Obligation: for each 3rd sample
Lentils	2	1600	x							x				x*				x	*Obligation: for each 3rd sample
Peas (chickpeas, chickling vetch)	1	2500	x							x				x*				x	*Obligation: for each 3rd sample
Other pulses, dry	2	1600	x							x									
4. OTHER SPROUTS AND SHOOTS																			
Other sprouts and shoots	1	2500	x							x									
5. CEREALS AND GRAIN FRUITS																			
Pseudocereals (Amaranth, Buckwheat, Chia seeds, Quinoa)	9	45	x							x						O*	x	x	*Recommendation: Copper
Corn (sweet corn)	1	2500	x							x									
Corn (popcorn corn)	1	2500	x							x									

Legend:

x

*

O

Multi methods:

GC

LC-MS/MS

2,4-D

¹The risk grouping of the products is made by QS and is based on a scale from 1 (lowest risk) to 9 (highest risk).

²At least the first sample and afterwards every 4th sample needs to be taken after all process steps

³For positive detections of Matrine by multi-method(s) a precise quantification of the active substance by a single method is required, if this is not ensured by the multi-methods.

⁴The first sample is required as soon as 10 percent or more of the tonnage defined in the control plan of a product has been purchased as QS produce within one year (approx. 12 months). If the product is purchased in smaller quantities as QS produce, no sampling is required.

Legend Risk groups - Tonnage

Risk groups	Tonnage calculation
1	2500
2	1600
3	875
4	640
5	480
6	340
7	170
8	90
9	45